arXiv:1609.09433v1 [cs.DS] 29 Sep 2016

Maximizing the Strong Triadic Closure in Split Graphs and Proper Interval Graphs

Athanasios Konstantinidis* Charis Papadopoulos†

Abstract

In social networks the Strong Triadic Closure is an assignment of the edges with strong or weak labels such that any two vertices that have a common neighbor with a strong edge are adjacent. The problem of maximizing the number of strong edges that satisfy the strong triadic closure was recently shown to be NP-complete for general graphs. Here we initiate the study of graph classes for which the problem is solvable. We show that the problem admits a polynomial-time algorithm for two unrelated classes of graphs: proper interval graphs and trivially-perfect graphs. To complement our result, we show that the problem remains NP-complete on split graphs, and consequently also on chordal graphs. Thus we contribute to define the first border between graph classes on which the problem is polynomially solvable and on which it remains NP-complete.

1 Introduction

Predicting the behavior of a network is an important concept in the field of social networks [6]. Understanding the strength and nature of social relationships has found an increasing usefulness in the last years due to the explosive growth of social networks (see e.g., [2]). Towards such a direction the Strong Triadic Closure principle enables to understand the structural properties of the underlying graph: it is not possible for two individuals to have a strong relationship with a common friend and not know each other [11]. Such a principle stipulates that if two people in a social network have a "strong friend" in common, then there is an increased likelihood that they will become friends themselves at some point in the future. Satisfying the Strong Triadic Closure is to characterize the edges of the underlying graph into weak and strong such that any two vertices that have a strong neighbour in common are adjacent. Since users interact and actively engage in social networks by creating strong relationships, it is natural to consider the MaxSTC problem: maximize the number of strong edges that satisfy the Strong Triadic Closure. The problem has been shown to be NP-complete for general graphs whereas its dual problem of minimizing the number of weak edges admits a constant factor approximation ratio [26].

In this work we initiate the computational complexity study of the MaxSTC problem in important classes of graphs. If the input graph is a P_3-free graph (i.e., a graph having no induced path on three vertices which is equivalent with a graph that consists of vertex-disjoint union of cliques) then there is a trivial solution by labeling strong all the edges. Such an observation might falsely lead into a graph modification problem, known as Cluster Deletion problem (see e.g., [12]), in which we want to remove the minimum number of edges that correspond to the weak edges, such that the resulting graph does not contain a P_3 as an induced subgraph. More precisely the obvious reduction would consist in labeling the deleted edges in the instance of Cluster Deletion as weak, and the remaining ones as

*Department of Mathematics, University of Ioannina, Greece. E-mail: konsakis@yahoo.com
†Department of Mathematics, University of Ioannina, Greece. E-mail: charis@cs.uoi.gr

strong. However, this reduction fails to be correct due to the fact that the graph obtained by deleting the weak edges in an optimal solution of MaxSTC may contain an induced P_3, so long as those three vertices induce a triangle in the original graph (prior to deleting the weak edges).

To the best of our knowledge, no previous results were known prior to our work when restricting the input graph for the MaxSTC problem. It is not difficult to see that for bipartite graphs the MaxSTC problem has a simple polynomial-time solution by considering a maximum matching that represent the strong edges [20]. It is well-known that a maximum matching of a graph corresponds to a maximum independent set of its line graph that represents the adjacencies between the edges [7]. As previously noted, for general graphs it is not necessarily the case that a maximum matching corresponds to the optimal solution for MaxSTC. Here we show a similar characterization for MaxSTC by considering the adjacencies between the edges of a graph that participate in induced P_3's. Such a characterization allows us to exhibit structural properties towards the computation of an optimal solution of MaxSTC.

Due to the nature of the P_3 existence that enforce the labeling of weak edges, there is an interesting connection to problems related to the *square root* of a graph; a graph H is a *square root* of a graph G and G is the *square* of H if two vertices are adjacent in G whenever they are at distance one or two in H. Any graph does not have a square root (for example consider a simple path), but every graph contains a subgraph that has a square root. Although it is NP-complete to determine if a given chordal graph has a square root [19], there are polynomial-time algorithms when the input is restricted to bipartite graphs [18], or proper interval graphs [19], or trivially-perfect graphs [23]. The relationship between MaxSTC and to that of determining square roots can be seen as follows. In the MaxSTC problem we are given a graph G and we want to select the maximum possible number of edges, at most one from each induced P_3 in G. Thus we need to find the largest subgraph (in terms of the number of its edges) H of G such that the square of H is a subgraph of G. However previous results related to square roots were concerned with deciding if the whole graph has a square root and there are no such equivalent formulations related to the largest square root.

Our main motivation is to understand the complexity of the problem on subclasses of chordal graphs, since the class of chordal graphs (i.e., graphs having no chordless cycle of length at least four) finds important applications in both theoretical and practical areas related to social networks [1, 16, 24]. More precisely two famous properties can be found in social networks. For most known social and biological networks their diameter, that is, the length of the longest shortest path between any two vertices of a graph, is known to be a small constant [14]. On the other hand it has been shown that the most prominent social network subgraphs are cliques, whereas highly infrequent induced subgraphs are cycles of length four [27]. Thus it is evident that subclasses of chordal graphs are close related to such networks, since they have rather small diameter (e.g., split graphs or trivially-perfect graphs) and are characterized by the absence of chordless cycles (e.g., proper interval graphs). Towards such a direction we show that MaxSTC is NP-complete on split graphs and consequently also on chordal graphs. On the positive side, we present the first polynomial-time algorithm for computing MaxSTC on proper interval graphs. Proper interval graphs, also known as unit interval graphs or indifference graphs, form a subclass of interval graphs and they are unrelated to split graphs. By our result they form the first graph class, other than bipartite graphs, for which MaxSTC is shown to be polynomial time solvable. In order to obtain our algorithm, we take advantage of their clique path and apply a dynamic programming on subproblems defined by passing the clique path in its natural ordering. Furthermore by considering the equivalent transformation of the problem mentioned earlier, we show

Figure 1: A split graph G is shown to the left side. The right side depicts a solution for MaxSTC on G where the weak edges are exactly the edges of G that are not shown.

that MaxSTC admits a simple polynomial-time solution on trivially-perfect graphs. Thus we contribute to define the first borderline between graph classes on which the problem is polynomially solvable and on which it remains NP-complete.

2 Preliminaries

All graphs considered here are simple and undirected. A graph is denoted by $G = (V, E)$ with vertex set V and edge set E. We use the convention that $n = |V|$ and $m = |E|$. The *neighborhood* of a vertex v of G is $N(v) = \{x \mid vx \in E\}$ and the *closed neighborhood* of v is $N[v] = N(v) \cup \{v\}$. For $S \subseteq V$, $N(S) = \bigcup_{v \in S} N(v) \setminus S$ and $N[S] = N(S) \cup S$. A graph H is a *subgraph* of G if $V(H) \subseteq V(G)$ and $E(H) \subseteq E(G)$. For $X \subseteq V(G)$, the subgraph of G *induced* by X, $G[X]$, has vertex set X, and for each vertex pair u, v from X, uv is an edge of $G[X]$ if and only if $u \neq v$ and uv is an edge of G. For $R \subseteq E(G)$, $G \setminus R$ denotes the graph $(V(G), E(G) \setminus R)$, that is a subgraph of G and for $S \subseteq V(G)$, $G - S$ denotes the graph $G[V(G) - S]$, that is an induced subgraph of G. Two adjacent vertices u and v are called *twins* if $N[u] = N[v]$.

A *clique* of G is a set of pairwise adjacent vertices of G, and a *maximal clique* of G is a clique of G that is not properly contained in any clique of G. An *independent set* of G is a set of pairwise non-adjacent vertices of G. For $k \geq 2$, the chordless path on k vertices is denoted by P_k and the chordless cycle on k vertices is denoted by C_k. For an induced path P_k, the vertices of degree one are called endvertices. A vertex v is *universal* in G if $N[v] = V(G)$ and v is *isolated* if $N(v) = \emptyset$. For two vertices u and v we say that u *sees* v if $\{u, v\} \in E(G)$; otherwise, we say that u *misses* v. We extend this notion to vertex sets: a set A sees (resp., misses) a vertex set B if every vertex of A is adjacent (resp., non-adjacent) to every vertex of B. We say that two *edges are non-adjacent* if they have no common endpoint; otherwise we call them *adjacent edges*.

Strong Triadic Closure. Given a graph $G = (V, E)$, a *strong-weak labeling* on the edges of G is a function λ that assigns to each edge of $E(G)$ one of the labels *strong* or *weak*; i.e., $\lambda : E(G) \to \{\text{strong}, \text{weak}\}$. An edge that is labeled strong (resp., weak) is simple called *strong* (resp. *weak*). The *strong triadic closure* of a graph G is a strong-weak labeling λ such that for any two strong edges $\{u, v\}$ and $\{v, w\}$ there is a (weak or strong) edge $\{u, w\}$. In other words, in a strong triadic closure there are no pair of strong edges $\{u, v\}$ and $\{v, w\}$ such that $\{u, w\} \notin E(G)$.

The problem of computing the maximum strong triadic closure, denoted by MaxSTC, is to find a strong-weak labeling on the edges of $E(G)$ that satisfies the strong triadic closure and has the maximum number of strong edges. Note that its dual problem asks for the minimum number of weak edges. Here we focus on maximizing the number of strong edges in a strong triadic closure.

Let G be a strong-weak labeled graph. We denote by (E_S, E_W) the partition of $E(G)$ into strong edges E_S and weak edges E_W. The graph spanned by E_S is the graph $G \setminus E_W$.

For a vertex $v \in V(G)$ we say that the *strong neighbour* of v is the other endpoint of a strong edge incident to v. We denote by $N_S(v) \subseteq N(v)$ the strong neighbours of v. Similarly we say that a vertex u is *strongly adjacent* to v if u is adjacent to v and the edge $\{u, v\}$ is strong.

Observation 2.1. *Let G be a strong-weak labeled graph. Then $G \setminus E_W$ satisfies the strong triadic closure if and only if for every P_3 in $G \setminus E_W$, the vertices of the P_3 induce a K_3 in G.*

Proof. Observe that $G \setminus E_W$ is the graph spanned by the strong edges. If for two strong edges $\{u, v\}$ and $\{v, w\}$, $\{u, w\} \notin E(G \setminus E_W)$ then $\{u, w\}$ is an edge in G and, thus, u, v, w induce a K_3 in G. On the other hand notice that any two strong edges of $G \setminus E_W$ are either non-adjacent or share a common vertex. If they share a common vertex then the vertices must induce a K_3 in G. □

Therefore in the MaxSTC problem we want to minimize the number of the removal (weak) edges E_W from G such that every three vertices that induce a P_3 in $G \setminus E_W$ form a clique in G. Then it is not difficult to see that $G \setminus E_W$ satisfies the strong triadic closure if and only if for every vertex v, $N_S[v]$ induces a clique in G.

3 MaxSTC on split graphs

Here we provide an NP-hardness result for MaxSTC on split graphs. A graph $G = (V, E)$ is a *split graph* if V can be partitioned into a clique C and an independent set I, where (C, I) is called a *split partition* of G. Split graphs form a subclass of the larger and widely known graph class of *chordal graphs*, which are the graphs that do not contain induced cycles of length 4 or more as induced subgraphs. It is known that split graphs are self-complementary, that is, the complement of a split graph remains a split graph. First we show the following result.

Lemma 3.1. *Let $G = (V, E)$ be a split graph with a split partition (C, I). Let E_S be the set of strong edges in an optimal solution for* MaxSTC *on G and let I_W be the vertices of I that are incident to at least one edge of E_S.*

1. *If every vertex of I_W misses at least three vertices of C then $E_S = E(C)$.*
2. *If every vertex of I_W misses exactly one vertex of C then $|E_S| \leq |E(C)| + \lfloor \frac{|I_W|}{2} \rfloor$.*

Proof. Let w_i be a vertex of I and let B_i be the set of vertices in C that are non-adjacent to w_i. Let A_i be the strong neighbors of w_i in an optimal solution. For the edges of the clique, there are $|A_i||B_i|$ weak edges due to the strong triadic closure. Moreover any vertex w_j of $I \setminus \{w_i\}$ cannot have a strong neighbor in A_i. This means that $A_i \cap A_j = \emptyset$. Notice, however, that both sets $B_i \cap B_j$ and $A_i \cap B_j$ are not necessarily empty.

Observe that I_W contains the vertices of I that are incident to at least one strong edge. Let $E(A, B)$ be the set of weak edges that have one endpoint in A_i and the other endpoint in B_i, for every $1 \leq i \leq |I_W|$. We show that $2|E(A, B)| \leq \sum_{w_i \in I_W} |A_i||B_i|$. Let $\{a, b\} \in E(A, B)$ such that $a \in A_i$ and $b \in B_i$. Assume that there is a pair A_j, B_j such that $\{a, b\}$ is an edge between A_j and B_j, for $j \neq i$. Then a cannot belong to A_j since $A_i \cap A_j = \emptyset$. Thus $a \in A_i$ and $b \in A_j$. Therefore for every edge $\{a, b\} \in E(A, B)$ there are at most two pairs (A_i, B_i) and (A_j, B_j) for which $a \in A_i \cup B_j$ and $b \in B_i \cup A_j$. This means that every edge of $E(A, B)$ is counted at most twice in $\sum_{w_i \in I_W} |A_i||B_i|$.

For any two edges $\{u, v\}, \{v, z\} \in E(C) \setminus E(A, B)$, observe that they satisfy the strong triadic closure since there is the edge $\{u, z\}$ in G. Thus the strong edges of the clique are exactly the set of edges $E(C) \setminus E(A, B)$. In total by counting the number of strong edges

between the independent set and the clique, we have $|E_S| = |E(C) \setminus E(A,B)| + \sum_{w_i \in I_W} |A_i|$. Since $2|E(A,B)| \leq \sum_{w_i \in I_W} |A_i||B_i|$, we get $|E_S| \leq |E(C)| + \sum_{w_i \in I_W} |A_i|(1 - \lfloor \frac{|B_i|}{2} \rfloor)$. Now the first claim of the lemma holds because $|B_i| = 3$ and $(1 - \lfloor \frac{|B_i|}{2} \rfloor) < 0$ so that $I_W = \emptyset$. For the second claim we show that for every vertex of I_W, $|A_i| = 1$. Let $w_i \in I_W$ such that $|A_i| \geq 2$ and let $B_i = \{b_i\}$. Recall that no other vertex of I_W has strong neighbours in A_i. Also note that there is at most one vertex w_j in I_W that has b_i as a strong neighbour. If such a vertex w_j exist and for the vertex b_j of the clique that misses w_j it holds $b_j \in A_i$, then we let $v = b_j$; otherwise we choose v as an arbitrary vertex of A_i. Observe that no vertex of $I \setminus \{w_i\}$ has a strong neighbour in $A_i \setminus \{v\}$ and only $w_j \in I_W$ is strongly adjacent to b_i. Then we label weak the $|A_i| - 1$ edges between w_i and the vertices of $A_i \setminus \{v\}$ and we label strong the $|A_i| - 1$ edges between b_i and the vertices of $A_i \setminus \{v\}$. Making strong the edges between b_i and the vertices of $A_i \setminus \{v\}$ does not violate the strong triadic closure since every vertex of $C \cup \{w_j\}$ is adjacent to every vertex of $A_i \setminus \{v\}$. Therefore for every vertex $w_i \in I_W$, $|A_i| = 1$ and by substituting $|B_i| = 1$ in the formula for $|E_S|$ we get the claimed bound. □

In order to give the reduction, we introduce the following problem that we call *maximum disjoint non-neighbourhood*: given a split graph (C, I) where every vertex of I misses three vertices from C, we want to find the maximum subset S_I of I such that the non-neighbourhoods of the vertices of S_I are pairwise disjoint. In the corresponding decision version, denoted by MaxDisjointNN, we are also given an integer k and the problem asks whether $|S_I| \geq k$. The polynomial-time reduction to MaxDisjointNN is given from the classical NP-complete problem 3-Set Packing [15]: given a universe $\mathcal{U}$ of n elements, a family $\mathcal{F}$ of triplets of $\mathcal{U}$, and an integer k, the problem asks for a subfamily $\mathcal{F}' \subseteq \mathcal{F}$ with $|\mathcal{F}'| \geq k$ such that all triplets of $\mathcal{F}'$ are pairwise disjoint.

Theorem 3.2. MaxDisjointNN *is NP-complete on split graphs.*

Proof. Given a split graph $G = (C, I)$ and $S_I \subseteq I$, checking whether S_I is a solution for MaxDisjointNN amounts to checking whether every pair of vertices of S_I have common neighbourhood. As this can be done in polynomial time the problem is in NP. We will give a polynomial-time reduction to MaxDisjointNN from the classical NP-complete problem 3-Set Packing [15]: given a universe $\mathcal{U}$ of n elements, a family $\mathcal{F}$ of triplets of $\mathcal{U}$, and an integer k, the problem asks for a subfamily $\mathcal{F}' \subseteq \mathcal{F}$ with $|\mathcal{F}'| \geq k$ such that all triplets of $\mathcal{F}'$ are pairwise disjoint.

Let $(\mathcal{U}, \mathcal{F}, k)$ be an instance of the 3-Set Packing. We construct a split graph $G = (C, I)$ as follows. The clique of G is formed by the n elements of $\mathcal{U}$. For every triplet F_i of $\mathcal{F}$ we add a vertex v_i in I that is adjacent to every vertex of C except the three vertices that correspond to the triplet F_i. Thus every vertex of I misses exactly three vertices from C and sees the rest of C. Now it is not difficult to see that there is a solution $\mathcal{F}'$ for 3-Set Packing$(\mathcal{U}, \mathcal{F}, k)$ of size at least k if and only if there is a solution S_I for MaxDisjointNN(G, k) of size at least k. For every pair (F_i, F_j) of $\mathcal{F}'$ we know that $F_i \cap F_j = \emptyset$ which implies that the vertices v_i and v_j have disjoint non-neighbourhood since F_i corresponds to the non-neighbourhood of v_i. By the one-to-one mapping between the sets of $\mathcal{F}$ and the vertices of I, every set F_i belongs to $\mathcal{F}'$ if and only if v_i belongs to S_I. □

Now we turn to our original problem MaxSTC. The decision version of MaxSTC takes as input a graph G and an integer k and asks whether there is strong-weak labeling of the edges of G that satisfies the strong triadic closure with at least k strong edges.

Theorem 3.3. *The decision version of* MaxSTC *is NP-complete on split graphs.*



Figure 2: The split graph $(C \cup C_Y, I \cup I_X)$ given in the polynomial-time reduction. Every vertex w_i misses the vertices of B_i and sees the vertices of $(C \cup C_Y) - B_i$. Every vertex x_i misses y_i and sees the vertices of $(C \cup C_Y) - \{y_i\}$. The sets $B_1, \ldots, B_k$ are pairwise disjoint whereas for every set B_j, $k < j \leq |I|$, there is a set B_i, $1 \leq i \leq k$, such that $B_i \cap B_j \neq \emptyset$. The drawn edges correspond to the strong edges between the independent set and the clique, and the dashed edges are the only weak edges in the clique $C \cup C_Y$.

Proof. Given a strong-weak labeling (E_S, E_W) of a split graph $G = (C, I)$, checking whether (E_S, E_W) satisfies the strong triadic closure amounts to check in $G - E_W$ whether there is a non-edge between its endvertices of every P_3 according to Observation 2.1. Thus by listing all P_3's of $G \setminus E_W$ the problem belongs to NP. Next we give a polynomial-time reduction to MaxSTC from the MaxDisjointNN problem on split graphs which is NP-complete by Theorem 3.2. Let (G, k) be an instance of MaxDisjointNN where $G = (C, I)$ is a split graph such that every vertex of the independent set I misses exactly three vertices from the clique C. For a vertex $w_i \in I$, we denote by B_i the set of the three vertices in C that are non-adjacent to w_i. Let $|C| = n$. We extend G and construct another split graph G' as follows (see Figure 2):

- We add n vertices $y_1, \ldots, y_n$ in the clique that consist the set C_Y.
- We add n vertices $x_1, \ldots, x_n$ in the independent set that consist the set I_X.
- For every $1 \leq i \leq n$, y_i is adjacent to all vertices of $(C \cup C_Y \cup I \cup I_X) \setminus \{x_i\}$.
- For every $1 \leq i \leq n$, x_i is adjacent to all vertices of $(C \cup C_Y) \setminus \{y_i\}$.

Thus w_i misses only the vertices of B_i from the clique. By construction it is clear that G' is a split graph with a split partition $(C \cup C_Y, I \cup I_X)$. Notice that the clique $C \cup C_Y$ has $2n$ vertices and $G = G'[I \cup C]$.

We claim that G has a solution for MaxDisjointNN of size at least k if and only if G' has a strong triadic closure with at least $n(2n-1) + \lfloor \frac{n}{2} \rfloor + \lceil \frac{k}{2} \rceil$ strong edges.

Assume that $\{w_1, \ldots, w_k\} \subseteq I$ is a solution for MaxDisjointNN on G of size at least k. Since the sets $B_1, \ldots, B_k$ are pairwise disjoint, there are k distinct vertices $y_1, \ldots, y_k$ in C_Y such that $k \leq n$. We will give a strong-weak labeling for the edges of G' that fulfills the strong triadic closure and has at least the claimed number of strong edges. For simplicity, we describe only the strong edges; the edges of G' that are not given are all labeled weak. We label the edges incident to each vertex w_i, y_i, x_i and the three vertices of each set B_i, for $1 \leq i \leq k$ as follows:

- The edges of the form $\{y_i, v\}$ are labeled strong if $v \in (C \cup C_Y) \setminus B_i$ or $v = w_i$.

- The edges incident to x_i and the three vertices of B_i are labeled strong.

Next we label the edges incident to the rest of the vertices. Let I_W be the vertices of $I \setminus \{w_1, \ldots, w_k\}$ and let C_W be the vertices of $C \setminus (B_1 \cup \cdots \cup B_k)$. No edge incident to a vertex of I_W is labeled strong. For every vertex $u \in C_W$ we label the edge $\{u, v\}$ strong if $v \in (C \cup C_Y)$. Let $C'_Y = \{y_{k+1}, \ldots, y_n\}$ and let $I'_X = \{x_{k+1}, \ldots, x_n\}$. Recall that every vertex x_{k+j} is adjacent to every vertex of $C'_Y \setminus \{y_{k+j}\}$. Let $\ell = \lfloor \frac{n-k}{2} \rfloor$. Let $M = \{e_1, \ldots, e_\ell\}$ be a maximal set of pairwise non-adjacent edges in $G'[C'_Y]$ where $e_j = \{y_{k+2j-1}, y_{k+2j}\}$, for $j \in \{1, \ldots, \ell\}$; note that M is a maximal matching of $G'[C'_Y]$. For every vertex $y \in C'_Y$, we label the edge $\{y, v\}$ strong if $v \in (C \cup C_Y) \setminus \{y'\}$ such that $\{y, y'\} \in M$. Moreover, for $j \in \{1, \ldots, \ell\}$, the edges $\{x_{k+2j-1}, y_{k+2j}\}$ and $\{x_{k+2j}, y_{k+2j-1}\}$ are labeled strong. Note that if $n - k$ is odd then no edge incident to the unique vertex y_n belongs to M and all edges between y_n and the vertices of $C \cup C_Y$ are labeled strong; in such a case also note that no edge incident to x_n is strong.

Let us show that such a labeling fulfills the strong triadic closure. Any labeling for the edges inside $G'[C \cup C_Y]$ is satisfied since $G'[C \cup C_Y]$ is a clique. Also note that there are no two adjacent strong edges that have a common endpoint in the clique $C \cup C_Y$ and the two other endpoints in the independent set $I \cup I_X$. If there are two strong edges incident to the same vertex v of the independent set then $v \in \{x_1, \ldots, x_k\}$ and $N_S[v] = B_i$ is a clique. Assume that there are two adjacent strong edges $\{u, v\}$ and $\{v, z\}$ such that $u \in I \cup I_X$, and $v, z \in C \cup C_Y$.

- If $u \in \{w_1, \ldots, w_k\}$ then $\{u, z\} \in E(G')$ since every w_i misses only the vertices of B_i.
- If $u \in \{x_1, \ldots, x_k\}$ then $v \in B_i$ and $\{u, z\} \in E(G')$ since every vertex x_i misses only y_i.
- If $u \in I_X \setminus \{x_1, \ldots, x_k\}$ then the strong neighbours of v in $C \cup C_Y$ are adjacent to u in G' since for the only non-neighbour of u in $C \cup C_Y$ there is a weak edge incident to v.

Recall that there is no strong edge incident to the vertices of $I \setminus \{w_1, \ldots, w_k\}$. Therefore the given strong-weak labeling fulfills the strong triadic closure.

Observe that the number of vertices in $C \cup C_Y$ is $2n$. There are exactly $3k + \ell$ weak edges in $G'[C \cup C_Y]$. Thus the number of strong edges in $G'[C \cup C_Y]$ are $n(2n-1) - 3k - \ell$. There are k strong edges incident to $\{w_1, \ldots, w_k\}$, $3k$ strong edges incident to $\{x_1, \ldots, x_k\}$, and 2ℓ strong edges incident to the vertices of $I_X \setminus \{x_1, \ldots, x_k\}$. Thus the total number of strong edges is $n(2n-1) - 3k - \ell + k + 3k + 2\ell = n(2n-1) + \ell + k$ and by substituting $\ell = \lfloor \frac{n-k}{2} \rfloor$ we get the claimed bound.

For the opposite direction, assume that G' has a strong triadic closure with at least $n(2n-1) + \lfloor \frac{n}{2} \rfloor + \lceil \frac{k}{2} \rceil$ strong edges. Let E_S be the set of strong edges in such a strong-weak labeling. Observe that the number of edges in $G'[C \cup C_Y]$ is $n(2n-1)$ which implies that E_S contains edges between the independent set $I \cup I_X$ and the clique $C \cup C_Y$. If no vertex of I_X is incident to an edge of E_S then the first statement of Lemma 3.1 implies that $|E_S| = |E(C \cup C_Y)| = n(2n-1)$. And if no vertex of I is incident to an edge of E_S then the second statement of Lemma 3.1 shows that $|E_S| \leq |E(C \cup C_Y)| + \lfloor \frac{n}{2} \rfloor$. Therefore E_S contains edges that are incident to a vertex of I and edges that are incident to a vertex of I_X.

In the graph spanned by E_S we denote by S_W the set of vertices of I that have strong neighbours in $C \cup C_Y$. We will show that the non-neighbourhoods of the vertices of S_W in $C \cup C_Y$ are disjoint in G' and, since G is an induced subgraph of G', their non-neighbourhoods are also disjoint in G.

Claim 3.4. *For every $w_i \in S_W$, $N_S(w_i) \subseteq C_Y$ and there exists a unique vertex $x \in I_X$ such that $N_S(x) = B_i$.*

Proof: Let w_i be a vertex of S_W. We first show that $N_S(w_i) \subseteq C_Y$. Let W_i be the strong neighbours of w_i in C and let Y_i be the strong neighbours of w_i in C_Y. Observe that no other vertex of S_W has a strong neighbour in $W_i \cup Y_i$. Further notice that there are $(|W_i|+|Y_i|)|B_i|$ weak edges since w_i is non-adjacent to the vertices of B_i. We show that for every vertex $w_i \in S_W$ it holds $W_i = \emptyset$. For all vertices w_i for which $W_i \neq \emptyset$ we replace in E_S the strong edges between w_i and the vertices of W_i by the edges between the vertices of B_i and W_i. Notice that making strong the edges between the vertices of B_i and W_i does not violate the strong triadic closure since no vertex from S_W has a strong neighbour in B_i and every vertex of I_X is adjacent to all the vertices of W_i. Since $\sum |W_i| < \sum |B_i||W_i|$, such a replacement is safe for the number of edges of E_S. Thus every vertex $w_i \in S_W$ has strong neighbours only in C_Y.

Let X_i be the set of vertices of I_X that have at least one non-neighbour in Y_i. By construction every vertex of Y_i is non-adjacent to exactly one vertex of I_X, and thus $|X_i| = |Y_i|$. Since w_i has strong neighbours in Y_i, every edge between X_i and Y_i is weak. By the previous argument every vertex of S_W has strong neighbours only in C_Y so that $N_S(B_i) \cap I = \emptyset$. Also notice that no two vertices of the independent set have a common strong neighbour in the clique, which means that there are at most $|B_i|$ strong neighbours between the vertices of B_i and I_X. Choose an arbitrary vertex $x \in X_i$. We replace all strong edges in E_S between B_i and I_X by $|B_i|$ strong edges between x and the vertices of B_i. Notice that such a replacement is safe since the unique non-neighbour of x belongs to Y_i and there are weak edges already in the solution between B_i and Y_i because of the strong edges between w_i and Y_i. Thus $B_i \subseteq N_S(x)$. We focus on the edges between the vertices of $(C \cup C_Y) \setminus (B_i \cup Y_i)$ and x. If a vertex x of X_i has a strong neighbour u in $(C \cup C_Y) \setminus B_i$ then the edge $\{u, y\}$ is weak where $y \in Y_i$ is the unique non-neighbour of x. Also notice that $N_S(u) \cap (I \cup I_X) = \{x\}$, $N_S(y) \cap (I \cup I_X) = \{w_i\}$, and w_i is adjacent to u. Then we can safely replace the strong edge $\{x, u\}$ by the edge $\{u, y\}$ and keep the same size of E_S. Hence $N_S(x) = B_i$. ♢

Claim 3.5. *For every $w_i \in S_W$, $N_S(w_i) = \{y\}$ where $y \in C_Y$ is the non-neighbour of x with $N_S(x) = B_i$.*

Proof: Let $Y_i = N_S(w_i)$. By Claim 3.4 we know that $Y_i \subseteq C_Y$ and there exists $x \in I_X$ such that $N_S(x) = B_i$. Both w_i and x are vertices of the independent set and, thus, no other vertex of $I \cup I_X$ has strong neighbours in $B_i \cup Y_i$. This means that if we remove w_i from S_W by making weak the edges incident to w_i and the vertices of Y_i then the edges between the vertices of B_i and $Y_i \setminus \{y\}$ are safely turned into strong. Let E'_S be the set of strong edges in an optimal solution such that all edges incident to w_i are weak. Then $|E_S| - |E'_S| = |Y_i| + |B_i| - |Y_i||B_i|$ and $|E_S| > |E'_S|$ only if $|Y_i| = 1$ because $|B_i| > 1$. Thus $N_S(w_i)$ contains exactly one vertex $y \in C_Y$. ♢

We claim that for every pair of vertices $w_i, w_j \in S_W$, $B_i \cap B_j = \emptyset$. Assume for contradiction that $B_i \cap B_j \neq \emptyset$. Applying Claim 3.4 for w_i shows that there exists $x \in I_X$ that has strong neighbours in every vertex of $B_i \cap B_j$. With a similar argument for w_j we deduce that there exists $x' \in I_X$ that has strong neighbours in every vertex of $B_i \cap B_j$. If $x \neq x'$ then a vertex from $B_i \cap B_j$ has two distinct strong neighbours in I_X which is not possible due to the strong triadic closure. Thus $x = x'$. Claim 3.5 implies that the unique non-neighbour y of x is strongly adjacent to both w_i and w_j. This however violates the strong triadic closure for the edges of E_S since w_i, w_j are non-adjacent and we reach a contradiction. Thus $B_i \cap B_j = \emptyset$. This means that the number of edges in E_S is at least $n(2n-1) + \lfloor \frac{n}{2} \rfloor + \lceil \frac{|S_W|}{2} \rceil$

which is maximized for $k = |S_W|$. Therefore E_S contains the maximum number of $|S_W|$ which is a solution for MaxDisjointNN on G, since G is an induced subgraph of G'. □

4 Computing MaxSTC on proper interval graphs

A graph is a *proper interval graph* if there is a bijection between its vertices and a family of closed intervals of the real line such that two vertices are adjacent if and only if the two corresponding intervals overlap and no interval is properly contained in another interval. A vertex ordering σ is a linear arrangement $\sigma = \langle v_1, \ldots, v_n \rangle$ of the vertices of G. For a vertex pair x, y we write $x \preceq y$ if $x = v_i$ and $y = v_j$ for some indices $i \leq j$; if $x \neq y$ which implies $i < j$ then we write $x \prec y$. The first position in σ will be referred to as the *left end* of σ, and the last position as the *right end*. We will use the expressions *to the left of*, *to the right of*, *leftmost*, and *rightmost* accordingly.

A vertex ordering σ for G is called a *proper interval ordering* if for every vertex triple x, y, z of G with $x \prec y \prec z$, $xz \in E(G)$ implies $xy, yz \in E(G)$. Proper interval graphs are characterized as the graphs that admit such orderings, that is, a graph is a proper interval graph if and only if it has a proper interval ordering [21]. We only consider this vertex ordering characterization for proper interval graphs. Moreover it can be decided in linear time whether a given graph is a proper interval graph, and if so, a proper interval ordering can be generated in linear time [21]. It is clear that a vertex ordering σ for G is a proper interval ordering if and only if the reverse of σ is a proper interval ordering. A connected proper interval graph without twin vertices has a unique proper interval ordering σ up to reversal [5, 13].

Let us turn our attention to the MaxSTC problem. Instead of maximizing the strong edges of the original graph G, we will look at the maximum independent set of the following graph that we call the *line-incompatibility* graph $\widehat{G}$ of G: for every edge of G there is a node in $\widehat{G}$ and two nodes of $\widehat{G}$ are adjacent if and only if the vertices of the corresponding edges induce a P_3 in G. Note that the line-incompatibility graph of G is a subgraph of the line graph[1] of G.

Proposition 4.1. *A subset S of edges $E(G)$ is an optimal solution for* MaxSTC *of G if and only if S is a maximum independent set of $\widehat{G}$.*

Proof. By Observation 2.1 for every P_3 in G at least one of its two edges must be labeled weak in S. This means that these two edges are adjacent in $\widehat{G}$ and they cannot belong to an independent set of $\widehat{G}$. On the other hand, by construction two nodes of $\widehat{G}$ are adjacent if and only if there is a P_3 in G. Thus the nodes of an independent set of $\widehat{G}$ can be labeled strong in G satisfying the strong triadic closure. □

Therefore we seek for the optimal solution of G by looking at a solution for a maximum independent set of $\widehat{G}$. We denote by $I_{\widehat{G}}$ the maximum independent set of $\widehat{G}$. To distinguish the vertices of $\widehat{G}$ with those of G we refer to the former as nodes and to the latter as vertices. For an edge $\{u, v\}$ of G we denote by uv the corresponding node of $\widehat{G}$.

Lemma 4.2. *Let x and y be twin vertices of a graph G. Then $xy \in I_{\widehat{G}}$ and for every vertex $u \in N(x)$, $xu \in I_{\widehat{G}}$ if and only if $yu \in I_{\widehat{G}}$.*

Proof. First we show that xy is an isolated vertex in $I_{\widehat{G}}$. If xy is adjacent to xu then y is non-adjacent to u in G which contradicts the fact that x and y are twins. Thus xy is

[1]The *line graph* of G is the graph having the edges of G as vertices and two vertices of the line graph are adjacent if and only if the two original edges are incident in G.

an isolated vertex in $\widehat{G}$ which implies $xy \in I_{\widehat{G}}$. For the second argument observe that for every vertex $u \in N(x)$, xu and yu are non-adjacent in $I_{\widehat{G}}$. Since $N[x] = N[y]$, nodes xu and yu have exactly the same neighborhood in $I_{\widehat{G}}$. This means xu is non-adjacent to a node $wz \in I_{\widehat{G}}$ if and only if yu is non-adjacent to wz. Thus if $xu \in I_{\widehat{G}}$ then yu is non-adjacent to any vertex of $I_{\widehat{G}}$ which shows that $yu \in I_{\widehat{G}}$, and if $xu \notin I_{\widehat{G}}$ then there is a node $wz \in I_{\widehat{G}}$ that is adjacent to both xu and yu. Hence $xu \in I_{\widehat{G}}$ if and only if $yu \in I_{\widehat{G}}$. □

Lemma 4.2 suggests to consider a graph G that has no twin vertices as follows. We partition $V(G)$ into sets of twins. A vertex that has no twin appears in its twin set alone. For every twin set W_x we choose an arbitrary vertex x and remove all its twin vertices except x from G. Let G' be the resulting graph that has no twin vertices. For every edge $\{x, y\}$ of G' we assign a weight equal to the product $|W_x| \cdot |W_y|$. This value corresponds to all edges of the original graph G between the vertices of W_x and W_y. The line-incompatibility graph $\widehat{G'}$ of G' is constructed as defined above with the only difference that a node of $\widehat{G'}$ has weight equal to the weight of its corresponding edge in G'. Let $I_{\widehat{G'}}$ be a *maximum weighted independent set*, that is an independent set of $\widehat{G'}$ such that the sum of the weights of its nodes is maximized. Then by Lemma 4.2 we have $I_{\widehat{G}} = I_{\widehat{G'}} \cup S(W)$ where $S(W)$ contains $|W_x|(|W_x| - 1)/2$ nodes for every twin set W_x. Therefore we are interested in computing the maximum weighted independent set of $\widehat{G'}$. Also note that G' is an induced subgraph of the original graph G. Since there is no ambiguity in the forthcoming results and in order to avoid heavier notation we refer to $\widehat{G'}$ as $\widehat{G}$ by assuming that G has no twin vertices and every vertex of G has a positive weight.

Let G be a proper interval graph and let σ be a proper interval ordering for G. We say that a solution $I_{\widehat{G}}$ has the *consecutive strong ordering* with respect to σ if the following holds for any three vertices x, y, z of G with $x \prec y \prec z$: $xz \in I_{\widehat{G}}$ implies $xy, yz \in I_{\widehat{G}}$. Our task is to show that such an optimal ordering exists. We start by characterizing the optimal solution $I_{\widehat{G}}$ with respect to the proper interval ordering σ.

Lemma 4.3. *Let x, y, z be three vertices of a proper interval graph G such that $x \prec y \prec z$. If $xz \in I_{\widehat{G}}$ then $xy \in I_{\widehat{G}}$ or $yz \in I_{\widehat{G}}$.*

Proof. We show that at least one of xy or yz belongs to $I_{\widehat{G}}$. Consider the node xy in $\widehat{G}$. If xy is adjacent to a node $xx_\ell \in I_{\widehat{G}}$ then $\{x_\ell, y\} \notin E(G)$. Then observe that $x_\ell \prec y$ because $x \prec y$ and $\{x_\ell, y\} \notin E(G)$. Since both xx_ℓ and xz belong to $I_{\widehat{G}}$, $\{x_\ell, z\} \in E(G)$. This however contradicts the proper interval ordering because $x_\ell \prec y \prec z$, $\{x_\ell, z\} \in E(G)$ and y is non-adjacent to x_ℓ. Thus xy is non-adjacent to any vertex $xx_\ell \in I_{\widehat{G}}$ and, in analogous fashion, yz is non-adjacent to any vertex $zz_r \in I_{\widehat{G}}$.

Now assume that xy is adjacent to a node $yy_r \in I_{\widehat{G}}$ and yz is adjacent to a node $y_\ell y \in I_{\widehat{G}}$. This means that $\{x, y_r\} \notin E(G)$ and $\{z, y_\ell\} \notin E(G)$. Since $\{x, z\} \in E(G)$, by the proper interval ordering we have $y_\ell \prec x \prec y \prec z \prec y_r$. Then notice that $\{y_\ell, y_r\} \in E(G)$, because both $yy_r, yy_\ell \in I_{\widehat{G}}$. By the proper interval ordering we know that both x and z are adjacent to y_ℓ, y_r, leading to a contradiction to the assumptions $\{x, y_r\} \notin E(G)$ and $\{z, y_\ell\} \notin E(G)$. Therefore at least one of xy or yz belongs to $I_{\widehat{G}}$. □

Thus by Lemma 4.3 we have two symmetric cases to consider. The next characterization suggests that there is a fourth vertex with important properties in each corresponding case.

Lemma 4.4. *Let x, y, z be three vertices of a proper interval graph G such that $x \prec y \prec z$ and $xz \in I_{\widehat{G}}$.*

- *If $xy \notin I_{\widehat{G}}$ and $yz \in I_{\widehat{G}}$ then xy is non-adjacent to any node $x_\ell x \in I_{\widehat{G}}$ and there is a vertex w such that $yw \in I_{\widehat{G}}$, $\{x, w\} \notin E(G)$, and $z \prec w$.*

- *If $xy \in I_{\widehat{G}}$ and $yz \notin I_{\widehat{G}}$ then yz is non-adjacent to any node $zz_r \in I_{\widehat{G}}$ and there is a vertex w such that $wy \in I_{\widehat{G}}$, $\{w, z\} \notin E(G)$ and $w \prec x$.*

Proof. Let $xy \notin I_{\widehat{G}}$ and $yz \in I_{\widehat{G}}$. The case for $xy \in I_{\widehat{G}}$ and $yz \notin I_{\widehat{G}}$ is completely symmetric. Assume for contradiction that there is no vertex w such that $yw \in I_{\widehat{G}}$, $\{x, w\} \notin E(G)$, and $z \prec w$. We prove that xy is non-adjacent to any node of $I_{\widehat{G}}$, contradicting the optimality of $I_{\widehat{G}}$. Suppose first that xy is adjacent to a node $x_\ell x \in I_{\widehat{G}}$. Then y is non-adjacent to x_ℓ in G. Notice that $x_\ell \prec x$ because y is adjacent to x and $x \prec y$. Due to the fact that $yz \in I_{\widehat{G}}$, we have that $x_\ell x$ and xz are non-adjacent in $\widehat{G}$ which implies that $\{x_\ell, z\} \in E(G)$. Since $x_\ell \prec x \prec y \prec z$ and $\{x_\ell, z\} \in E(G)$, by the proper interval ordering we get $\{x_\ell, y\} \in E(G)$ leading to a contradiction. Thus xy is non-adjacent to any node $x_\ell x \in I_{\widehat{G}}$.

Next assume that xy is adjacent to a node $yy_r \in I_{\widehat{G}}$. Then $\{x, y_r\} \notin E(G)$. By the assumption that there is no vertex w with $yw \in I_{\widehat{G}}$, $\{x, w\} \notin E(G)$, and $z \prec w$, we have $y_r \prec z$. This particularly means that $y_r \prec x$ or $x \prec y_r \prec z$. However both cases lead to a contradiction to $\{x, y_r\} \notin E(G)$ since in the former case we have $\{y_r, y\} \in E(G)$ and $y_r \prec x \prec y$, and in the latter case we know that $\{x, z\} \in E(G)$. Therefore xy has no neighbor in $I_{\widehat{G}}$ reaching a contradiction to the optimality of $I_{\widehat{G}}$. □

Now we are ready to show that that there is an optimal solution that has the described properties with respect to the given proper interval ordering.

Lemma 4.5. *There exists an optimal solution $I_{\widehat{G}}$ that has the consecutive strong property with respect to σ.*

Proof. Let σ be a proper interval ordering for G. Assume for contradiction that $I_{\widehat{G}}$ does not have the consecutive strong property. Then there exists a *conflict* with respect to σ, that is, there are three vertices x, y, z with $x \prec y \prec z$ and $xz \in I_{\widehat{G}}$ such that $xy \notin I_{\widehat{G}}$ or $yz \notin I_{\widehat{G}}$. We will show that as long as there are conflicts in σ, we can reduce the number of conflicts in σ without affecting the value of the optimal solution $I_{\widehat{G}}$. Consider such a conflict formed by the three vertices $x \prec y \prec z$ with $xz \in I_{\widehat{G}}$. By Lemma 4.3 we know that $xy \in I_{\widehat{G}}$ or $yz \in I_{\widehat{G}}$. Assume that $yz \in I_{\widehat{G}}$. Then clearly $xy \notin I_{\widehat{G}}$, for otherwise there is no conflict. Then by Lemma 4.4 there is a vertex w such that $yw \in I_{\widehat{G}}$, $\{x, w\} \notin E(G)$, and $x \prec y \prec z \prec w$. Notice that both triples x, y, z and y, z, w create conflicts in σ.

We start by choosing an appropriate such conflict that is formed by four vertices x, y, z, w so that $x \prec y \prec z \prec w$, $xz, yz, yw \in I_{\widehat{G}}$, and $\{x, w\} \notin E(G)$. Fix y and z in σ with y, z being the leftmost and the rightmost vertices, respectively, such that for every vertex v with $y \prec v \prec z$, $yv, vz \in I_{\widehat{G}}$ holds. Recall that $yz \in I_{\widehat{G}}$. We choose x as the leftmost vertex such that $xz \in I_{\widehat{G}}$ and we choose w as the rightmost vertex such that $yw \in I_{\widehat{G}}$. Observe that $\{x, w\} \notin E(G)$ since y and z participate in a conflict. Due to the properties of the considered conflicts all such vertices exist (see for e.g., Figure 3).

Let $W(x)$ be the set of vertices w_i such that $yw_i \in I_{\widehat{G}}$ and $\{x, w_i\} \notin E(G)$, and let $X(w)$ be the set of vertices x_j such that $x_j z \in I_{\widehat{G}}$ and $\{x_j, w\} \notin E(G)$. Observe that $w \in W(x)$ and $x \in X(w)$ such that w is the rightmost vertex in $W(x)$ and x is the leftmost vertex in $X(w)$. For a vertex w_i of $W(x)$ observe the following. If $w_i \prec x$ then $\{w_i, x\} \in E(G)$ because $\{w_i, y\} \in E(G)$ and if $x \prec w_i \prec z$ then $\{w_i, x\} \in E(G)$ because $\{x, z\} \in E(G)$. Thus $z \prec w_i$ which implies that $\{z, w_i\} \in E(G)$ since $\{y, w_i\} \in E(G)$. If $zw_i \in I_{\widehat{G}}$ then by the fact that $xz \in I_{\widehat{G}}$ we have $\{x, w_i\} \in E(G)$ contradicting the definition of $W(x)$. Moreover for every vertex b_1 such that $w_i b_1 \in I_{\widehat{G}}$ notice that $x \prec b_1$ since $\{x, w_i\} \notin E(G)$. If $x \prec b_1 \prec w_i$ then $\{z, b_1\} \in E(G)$ since $x \prec z \prec w_i$ and if $w_i \prec b_1$ then due to the fact that $yw_i, w_i b_1 \in I_{\widehat{G}}$ and $\{y, b_1\} \in E(G)$ we have again $\{z, b_1\} \in E(G)$, as $y \prec z \prec b_1$. Furthermore consider a vertex b_2 such that $z \prec b_2 \prec w$ and $b_2 \notin W(x)$. This means that $yb_2 \notin I_{\widehat{G}}$ or $yb_2 \in I_{\widehat{G}}$ with $\{b_2, x\} \in E(G)$. The latter case implies that b_2 is adjacent to every vertex of $X(w)$, since x

is the leftmost vertex in $X(w)$ and every vertex of $X(w)$ is to the left of z. Hence for every vertex w_i of $W(x)$ the following hold:

- $z \prec w_i$,
- $zw_i \notin I_{\widehat{G}}$,
- for every node $w_ib_1 \in I_{\widehat{G}}$, $\{z, b_1\} \in E(G)$, and
- for every vertex b_2 with $z \prec b_2 \prec w$ and $b_2 \notin W(x)$, $yb_2 \notin I_{\widehat{G}}$ or b_2 is adjacent to every vertex of $X(w)$.

With symmetric arguments for every vertex x_j of $X(w)$ we have the following:

- $x_j \prec y$,
- $x_jy \notin I_{\widehat{G}}$,
- for every node $a_1x_j \in I_{\widehat{G}}$, $\{a_1, y\} \in E(G)$, and
- for every vertex a_2 with $x \prec a_2 \prec y$ and $a_2 \notin X(w)$, $a_2z \notin I_{\widehat{G}}$ or a_2 is adjacent to every vertex of $W(x)$.

The topmost ordering given in Figure 3 illustrates the corresponding cases.

Let Y_w be the set of nodes yw_i in $\widehat{G}$ such that $w_i \in W(x)$, and let Z_x be the set of nodes x_jz in $\widehat{G}$ such that $x_j \in X(w)$. Observe that $Y_w, Z_x \subseteq I_{\widehat{G}}$ by the previous arguments. We show that removing either Y_w or Z_x from $I_{\widehat{G}}$ does not create any new conflict. Let $yw_i \in Y_w$ and let u be a vertex such that $uy \in I_{\widehat{G}}$ and $uw_i \in I_{\widehat{G}}$. If $y \prec u \prec w_i$ then no conflict is created by removing yw_i from $I_{\widehat{G}}$. Assume that $u \prec y \prec w_i$. Observe that $x \prec u \prec z$. Then $xu \notin I_{\widehat{G}}$ because $\{x, w_i\} \notin E(G)$. Since $xz \in I_{\widehat{G}}$ and at least one of xu, uz belongs to $I_{\widehat{G}}$, we have $uz \in I_{\widehat{G}}$. However this contradicts the leftmost choice for y in $x \prec u \prec y \prec z$ and there is no such vertex u. Next assume that $y \prec w_i \prec u$. Since w_i is non-adjacent to x and $w_i \prec u$, u is non-adjacent to x, as well. Then according to the definition of $W(x)$, $u \in W(x)$ and $yu \in Y_w$. The case for the nodes of Z_x is completely symmetric. Thus no conflicts are created by removing the nodes of Y_w or the nodes of Z_x from $I_{\widehat{G}}$.

Let Y_x be the set of nodes x_jy in $\widehat{G}$ such that $x_j \in X(w)$, and let Z_w be the set of nodes zw_i in $\widehat{G}$ such that $w_i \in W(x)$. We denote by $I(Y_x)$ and $I(Z_w)$ the following sets of nodes: $I(Y_x) = \left(I_{\widehat{G}} \setminus Y_w\right) \cup Y_x$ and $I(Z_w) = \left(I_{\widehat{G}} \setminus Z_x\right) \cup Z_w$. We show that both sets form independent sets in $\widehat{G}$. Consider the case for $I(Y_x)$. Let x_jy be a node of $I(Y_x)$. There are two cases to consider: there is a node $ux_j \in I_{\widehat{G}} \setminus Y_w \cup Y_x$ and $\{u, y\} \notin E(G)$ or there is a node $yv \in I_{\widehat{G}} \setminus Y_w \cup Y_x$ and $\{x_j, v\} \notin E(G)$. In the former case we know that $ux_j \in I_{\widehat{G}}$ and for every node $ux_j \in I_{\widehat{G}}$, $\{u, y\}$ must be an edge of G which leads to a contradiction to the non-adjacency of u and y. For the latter case observe that $yv \in I_{\widehat{G}} \setminus Y_w$ and $v \notin W(x)$. Since $\{x_j, v\} \notin E(G)$ and $\{y, v\} \in E(G)$, we have $z \prec v$ and by the rightmost choice of w for y we have $z \prec v \prec w$. This however implies that $z \prec v \prec w$, $v \notin W(x)$ and $yv \in I_{\widehat{G}}$ showing that $\{x_j, v\} \in E(G)$. Completely symmetric arguments hold for $I(Z_w)$. The two lowest orderings given in Figure 3 illustrate the considered cases. Thus $I(Y_x)$ and $I(Z_w)$ form independent sets in $\widehat{G}$.

Now observe that both $I(Y_x)$ and $I(Z_w)$ have a smaller number of conflicts with respect to $I_{\widehat{G}}$ because either x, y, z in $I(Y_x)$ or y, z, w in $I(Z_w)$ satisfy the consecutive strong property. It is clear that the difference between $I(Y_x)$ and $I_{\widehat{G}}$ are the nodes of Y_x and Y_w, whereas the difference between $I(Z_w)$ and $I_{\widehat{G}}$ are the nodes of Z_w and Z_x. For a set A of vertices having positive weights, denote by $M(A)$ the sum of the weights of its vertices. If $M(X(w)) \geq M(Z(x))$ then $M(I(Y_x)) \geq M(I_{\widehat{G}})$ and if $M(X(w)) < M(Z(x))$ then $M(I(Z_w)) > M(I_{\widehat{G}})$.



Figure 3: A proper interval ordering for a graph G with three different solutions considered in the proof of Lemma 4.5. A solid edge corresponds to a node of $\widehat{G}$ that belongs to $I_{\widehat{G}}$, which means that such an edge is labeled strong in an optimal strong-weak labeling, whereas a dashed edge corresponds to a node of $\widehat{G}$ that does not belong to $I_{\widehat{G}}$, which means that such an edge is labeled weak in an optimal strong-weak labeling. Observe that the lowest two orderings contain less *conflicts* than the topmost, that is, triple of vertices that violate the consecutive strong property.

Thus in any case we can replace appropriate set of nodes in $I_{\widehat{G}}$ and obtain an optimal solution with a smaller number of conflicts. Therefore by applying such a replacement in every such conflict, we get an optimal solution that has no conflicts and, thus, it satisfies the consecutive strong property. □

Lemma 4.5 suggests to find an optimal solution that has the consecutive strong property with respect to σ. In fact by Proposition 4.1 and the proper interval ordering, this reduces to computing the largest proper interval subgraph H of G such that the vertices of every P_3 of H induce a clique in G.

Let G be a proper interval graph and let $\sigma = \langle v_1, \ldots, v_n \rangle$ be its proper interval ordering. For a vertex v_i we denote by $v_{\ell(i)}$ and $v_{r(i)}$ its leftmost and rightmost neighbors, respectively, in σ. Observe that for any two vertices $v_i \prec v_j$ in σ, $v_{\ell(i)} \preceq v_{\ell(j)}$ and $v_{r(i)} \preceq v_{r(j)}$ [5]. For $1 \leq i \leq r(1)$, let $V_i = \{v_1, \ldots, v_i\}$, that is, V_i contains the *first* i vertices in σ. Given the set V_i, let r be an integer such that $1 \leq i \leq r \leq r(1)$.

Let $A(G)$ be the value of an optimal solution $I_{\widehat{G}}$ for G. For the set V_i we denote by $B(V_i)$ the value that corresponds to the total weight of the edges incident to v_1 and each of $v_2, \ldots, v_i$. Observe that any subset of vertices of V_i induces a clique in G. We denote

by $C(V_i)$ the value that corresponds to the total weight of the edges among all vertices of V_i. Given the first vertices V_i and $i \leq r \leq r(1)$, let $A(G, V_i, r)$ be the value of the optimal solution $I(G, V_i, r)$ in G such that every edge among the vertices of V_i belongs to $I(G, V_i, r)$ and $uv_k \notin I(G, V_i, r)$ with $u \in V_i$ and $k > r$.

Lemma 4.6. *Let G be a proper interval graph and let V_i and r such that $1 \leq i \leq r \leq r(1)$. Then, $A(G) = A(G, \{v_1\}, r(1))$, and*

$$A(G, V_i, r) = \begin{cases} \max\limits_{i \leq j \leq r} \{A(G - \{v_1\}, V_j \setminus \{v_1\}, r) + B(V_j)\} & \text{if } i < r, \\ A(G - V_i, \{v_{i+1}\}, r(i+1)) + C(V_i) & \text{if } i = r \text{ and } i < n, \\ C(V_i) & \text{otherwise.} \end{cases}$$

Proof. We show that $A(G)$ computes the value of an optimal solution that satisfies the consecutive strong property. By Lemma 4.5 such an ordering exists. Since there is no edge between v_1 and v_k with $k > r(1)$, it follows that $A(G) = A(G, \{v_1\}, r(1))$. Observe that every induced subgraph of a proper interval graph is proper interval, which implies that the graphs $G - \{v_1\}$ and $G - V_i$ remain proper interval. Let $V_i = \{v_1, \ldots, v_i\}$ be the first i vertices of G. According to Lemma 4.5 if $v_i v_k \notin I(G, V_i, r)$ with $k > i$, then every node $v_j v_k$, $1 \leq j \leq i$, does not belong to $I(G, V_i, r)$. Again by Lemma 4.5 if $v_1 v_k \in I(G, V_i, r)$ with $k > i$, then every node $v_j v_k$, $1 \leq j \leq i$, belongs to $I(G, V_i, r)$.

Let $I(V_i)$ be the nodes of $\widehat{G}$ corresponding to the edges of $G[V_i]$. Observe that by the definition of $I(G, V_i, r)$, $I(V_i) \subseteq I(G, V_i, r)$. If $i = r$ then it is clear that the nodes of $I(V_i)$ belong to $I(G, V_i, r)$. Moreover if $i = r < n$ then no node $v_i v_k$, with $i < k$, can be added in $I(G, V_i, r)$, implying that all the nodes corresponding to the edges between V_i and $G - V_i$ do not belong to $I(G, V_i, r)$. Thus we can safely add $I(V_i)$ to an optimal solution for $G - V_i$, that is, $I(G - V_i, \{v_{i+1}\}, r(i+1))$.

Assume that $i < r$. To see that $I(V_i) \subseteq I(G, V_i, r)$, notice that $V_i \subseteq V_j$ for every $i \leq j \leq r$ and $C(V_i) = B(V_i) + B(V_i \setminus \{v_1\}) + \cdots + B(\{v_{i-1}, v_i\})$. Consider the nodes $v_1 v_j$ of $\widehat{G}$ corresponding to the edges incident to v_1. Let $v_1 v_j \in I_{\widehat{G}}$ with j as large as possible. Then $i \leq j \leq r \leq r(1)$. By Lemma 4.5 every node $v_1 v_{j'}$ of $\widehat{G}$ with $1 \leq j' \leq j$ belongs to $I_{\widehat{G}}$. Thus $I(V_j) \subseteq I(G, V_i, r)$. Furthermore for every node $v_j v_k$, with $r \leq r(1) < k$, we know that $v_j v_k$ is adjacent to $v_1 v_j$ in $\widehat{G}$ since $\{v_1, v_k\} \notin E(G)$. Thus $v_j v_k \notin I_{\widehat{G}}$ which implies that every node $v_{j'} v_k$ with $j' \leq j$ does not belong to $I_{\widehat{G}}$. This particularly means that $v_j v_k \in I(G, V_i, r)$ only if $k \leq r \leq r(1)$. Therefore given an optimal solution by removing v_1 for all such possible values for j, we obtain the desired solution. □

Now we are equipped with our necessary tools in order to obtain our main result, namely a polynomial-time algorithm that solves the MaxSTC problem on a proper interval graph G.

Theorem 4.7. *There is a polynomial-time algorithm that computes the* MaxSTC *of a proper interval graph.*

Proof. Let G be a proper interval graph on n vertices and m edges. We first compute its proper interval ordering σ in linear time [21]. Then we compute its twin sets by using the fact that u and v are twins if and only if $\ell(u) = \ell(v)$ and $r(u) = r(v)$. Contracting the twin sets according to Lemma 4.2 results in a proper interval graph in which every vertex is associated with a positive weight. In order to compute the optimal solution $A(G)$ we use a dynamic programming approach based on its recursive formulation given in Lemma 4.6. Correctness follows from Proposition 4.1 and Lemmata 4.5 and 4.6.

Regarding the running time, notice that given the ordering σ we can remove the twin vertices in linear time. All instances of $A(G, V_i, r)$ can be computed as follows. Given the first

vertex v_1 we compute all possible sets V_i which are bounded by n. Since $1 \leq r \leq r(1)$, the number of instances $A(G, V_i, r)$ generated by v_1 is at most n^2. Also observe that computing the values $B(V_i)$ and $C(V_i)$ takes $O(m)$ time. Therefore the total running time of the algorithm is $O(n^3 m)$. □

5 Concluding remarks

Given the first study with positive and negative results for the MaxSTC problem on restricted input, there are some interesting open problems. Despite the structural properties that we proved for the solution on proper interval graphs, the complexity of MaxSTC on interval graphs is still open. Determining the complexity of MaxSTC for other graph classes towards AT-free graphs seems interesting direction for future work. More precisely, by Proposition 4.1 it is interesting to consider the line-incompatibility graph of comparability graphs since they admit a well-known similar characterization [17]. Moreover it is natural to characterize the graphs for which their line-incompatibility graph is perfect. Such a characterization will lead to to further polynomial cases of MaxSTC, since the maximum independent set of perfect graphs admits a polynomial solution [10]. A typical example is the class of bipartite graphs for which their line graph coincides with their line-incompatibility graph and it is known that the line graph of a bipartite graph is perfect (see for e.g., [3]). As we show next, another paradigm of this type is the class of trivially-perfect graphs.

A graph G is called *trivially-perfect* (also known as *quasi-threshold*) if for each induced subgraph H of G, the number of maximal cliques of H is equal to the maximum size of an independent set of H. It is known that the class of trivially-perfect graphs coincides with the class of (P_4, C_4)-free graphs, that is every trivially-perfect graph has no induced P_4 or C_4 [9]. A *cograph* is a graph without an induced P_4, that is a cograph is a P_4-free graph. Hence trivially-perfect graphs form a subclass of cographs.

Theorem 5.1. *The line-incompatibility graph of a trivially-perfect graph is cograph.*

Proof. Let G be a trivially-perfect graph, that is G is a (P_4, C_4)-free graph. We will show that the line-incompatibility graph $\widehat{G}$ of G is a P_4-free graph. Consider any P_3 in $\widehat{G}$. Due to the construction of $\widehat{G}$, the P_3 has one of the following forms: (i) v_1v_2, v_2v_3, v_3v_4 or (ii) v_1x, v_2x, v_3x. We prove that the P_3 has the second form because G has no induced P_4 or C_4. If (i) applies then $\{v_1, v_3\}, \{v_2, v_4\} \notin E(G)$ and $\{v_1, v_2\}, \{v_2, v_3\}, \{v_3, v_4\} \in E(G)$ which implies that $v_4 \neq v_1$. Thus G contains a P_4 or a C_4 depending on whether there is the edge $\{v_1, v_4\}$ in G. Hence every P_3 in $\widehat{G}$ has the form v_1x, v_2x, v_3x where v_1, v_2, v_3, x are distinct vertices of G. Now assume for contradiction that $\widehat{G}$ contains a P_4. Then the P_4 is of the form v_1x, v_2x, v_3x, v_4x because it contains two induced P_3's. The structure of the P_4 implies that $\{v_1, v_2\}, \{v_2, v_3\}, \{v_3, v_4\} \notin E(G)$ and $\{v_1, v_3\}, \{v_2, v_4\}, \{v_4, v_1\} \in E(G)$. This however shows that the vertices v_3, v_1, v_4, v_2 induce a P_4 in G leading to a contradiction that G is a (P_4, C_4)-free graph. Therefore $\widehat{G}$ is a P_4-free graph. □

By Theorem 5.1 and the fact that the maximum independent set of a cograph can be computed in linear time [4], MaxSTC can be solved in polynomial time on trivially-perfect graphs. We would like to note that the line-incompatibility graph of a cograph or a proper interval graph is not necessarily a perfect graph.

More general there are extensions and variations of the MaxSTC problem that are interesting to consider as proposed in [26]. A natural graph modification problem is the problem of adding the minimum number of edges that satisfy the strong triadic closure having as few as possible weak edges in the original graph. More formally the objective is to add F edges in G so that the resulting graph satisfies the strong triadic closure with the minimum number of $|F| + |E_W|$.

References

[1] A. B. Adcock, B. D. Sullivan, M. W. Mahoney. Tree decompositions and social graphs. *Internet Mathematics*, 12:315–361, 2016.

[2] L. Backstrom and J. Kleinberg. Romantic partnerships and the dispersion of social ties: a network analysis of relationship status on facebook. In *Proceedings of CSCW 2014*, pp. 831–841, 2014.

[3] A. Brandstädt, V.B. Le, and J. Spinrad. *Graph Classes – a Survey*. SIAM Monographs in Discrete Mathematics and Applications, SIAM, Philadelphia, 1999.

[4] D.G. Corneil, Y. Perl, and L.K. Stewart. Complement reducible graphs. *Discrete Applied Mathematics*, 3:163–174, 1981.

[5] X. Deng, P. Hell, and J. Huang. Linear-time representation algorithms for proper circular-arc graphs and proper interval graphs. *SIAM J. Comput.*, 25:390–403, 1996.

[6] D. Easley and J. Kleinberg. *Networks, Crowds, and Markets: Reasoning About a Highly Connected World*. Cambridge University Press, 2010.

[7] J. Edmonds. Paths, trees and flowers. *Canad. J. Math.*, 17:449–467, 1965.

[8] D. R. Fulkerson and O. A. Gross. Incidence matrices and interval graphs. *Pacific Journal of Mathematics*, 15:835–855, 1965.

[9] M.C. Golumbic. Trivially perfect graphs. *Discrete Mathematics*, 24:105-107, 1978.

[10] M. Götschel, L. Lovász, and A. Scrijver. Polynomial algorithms for perfect graphs. *Annals of Discrete Mathematics*, 21:325–356, 1984.

[11] M. Granovetter. The strength of weak ties. *American Journal of Sociology*, 78:1360–1380, 1973.

[12] P. Heggernes, D. Lokshtanov, J. Nederlof, C. Paul, and J. A. Telle. Generalized graph clustering: recognizing (p, q)-cluster graphs. In *Proceedings of WG 2010*, pp. 171–183, 2010.

[13] L. Ibarra. The clique-separator graph for chordal graphs. *Discrete Applied Mathematics*, 157:1737-1749, 2009.

[14] M. O. Jackson. *Social and economic networks*. Princeton university press, vol. 3, 2008.

[15] R. M. Karp. Reducibility among combinatorial problems. *Complexity of Computer Computations*, 85–103, 1972.

[16] D. J. Kleitman and R. V. Vohra. Computing the bandwidth of interval graphs. *SIAM J. Disc. Math.*, 3:373–375, 1990.

[17] D. Kratsch, R. M. McConnell, K. Mehlhorn, and J. P. Spinrad. Certifying algorithms for recognizing interval graphs and permutation graphs. *SIAM J. Comput.*, 36: 326–353, 2006.

[18] L. C. Lau. Bipartite roots of graphs. *ACM Transactions on Algorithms*, 2:178–208, 2006.

[19] L. C. Lau and D. G. Corneil. Recognizing powers of proper interval, split, and chordal graphs. *SIAM J. Disc. Math.*, 18: 83–102, 2004.

[20] F.T. Leighton and S. Rao. An approximate max-flow min-cut theorem for uniform multicommodity flow problems with applications to approximation algorithms. In *Proceedings of FOCS 1988*, pp. 422–431, 1998.

[21] P. J. Looges and S. Olariu. Optimal greedy algorithms for indifference graphs. *Computers & Mathematics with Applications*, 25:1525, 1993.

[22] N. Mahadev and U. Peled. *Threshold graphs and related topics.* Annals of Discrete Mathematics 56. North Holland, 1995.

[23] M. Milanič and O. Schaudt. Computing square roots of trivially perfect and threshold graphs. *Discrete Applied Mathematics*, 161:1538-1545, 2013.

[24] J. L. Pfaltz. Chordless cycles in networks. In *Proceedings of ICDE Workshops*, pp. 223–228, 2013.

[25] F. S. Roberts. Indifference Graphs. In *Proof Techniques in Graph Theory*, Academic Press, New York, 139–146, 1969.

[26] S. Sintos and P. Tsaparas. Using strong triadic closure to characterize ties in social networks, In *Proceedings of KDD 2014*, pp. 1466–1475, 2014.

[27] J. Ugander, L. Backstrom, and J. Kleinberg. Subgraph frequencies: Mapping the empirical and extremal geography of large graph collections. In *Proceedings of WWW 2013*, pp. 1307–1318, 2013.